

Supplementary Table S3. Detailed data for companion drugs used with amikacin liposome inhalation suspension in patients with refractory *Mycobacterium avium* complex pulmonary disease

Patient*	Antibiotics and duration, mo					
	Macrolide [†]	Ethambutol	Rifamycin	Moxifloxacin	Clofazimine	Prothionamide
1	Yes	Yes	Yes	Yes	-	-
	13.3	13.3	13.3	0.9	-	-
2	Yes	Yes	-	Yes	Yes	Yes
	11.7	3.6	-	7.5	0.8	0.7
3	Yes	Yes	-	Yes	Yes	-
	7.1	7.1	-	1.1	6.0	-
4	Yes	Yes	-	Yes	-	-
	10.3	10.3	-	10.3	-	-
5	Yes	Yes	-	-	Yes	-
	2.8	2.8	-	-	2.8	-
6	Yes	Yes	Yes	Yes	Yes	-
	7.1	9.8	2.7	7.1	5.7	-
Overall duration, mo	8.7 (7.1–11.4)	8.5 (4.5–10.2)	8.0 (2.7–13.3)	7.1 (1.1–7.5)	4.2 (2.3–5.8)	0.7

Values are presented as median (interquartile range).

*Patient no. 1, 2, and 5 had *M. avium*-disease; no. 3 and 4 had *Mycobacterium intracellulare*-disease. Patient no. 6 had co-infection with *M. avium* and *M. intracellulare*. [†]Azithromycin or clarithromycin.

-: negative.