

**Supplementary Table S4.** Significant factors for recent smoking cessation after policy change (2015) among men smokers

	Univariate analysis			Multivariate analysis		
	OR	95% CI	p-value	OR	95% CI	p-value
<b>Age, yr</b>						
19–29	1.82	0.68–4.83	0.229	1.77	0.66–4.77	0.259
30–39	2.51	0.91–6.95	0.076	1.78	0.62–5.15	0.283
40–49	1			1		
50–59	1.71	0.64–4.59	0.287	1.92	0.71–5.22	0.200
≥60	2.67	1.12–6.38	0.026	2.82	1.13–7.02	0.026
BMI	1.02	0.95–1.01	0.636			
Diabetes	1.39	0.64–3.02	0.402			
Hypertension	0.56	0.32–0.97	0.038	0.46	0.25–0.81	0.008
Married	1.82	0.92–3.57	0.084			
<b>Education level</b>						
Below elementary	2.28	1.10–4.73	0.027	2.11	0.96–4.67	0.064
Middle school	0.95	0.38–2.35	0.907	0.81	0.33–1.97	0.635
High school	1					
Above university	1.94	1.04–3.63	0.038	2.37	1.18–4.77	0.016
<b>Type of residence</b>						
General type	1					
Apartment	0.57	0.33–1.00	0.051			
<b>Household income</b>						
Lowest quartile (Q1)	1.16	0.51–2.64	0.731	1.40	0.60–3.29	0.435
Low–middle quartile (Q2)	2.32	1.16–4.64	0.018	2.66	1.24–5.72	0.013
High–middle quartile (Q3)	0.94	0.46–1.92	0.873	0.87	0.39–1.94	0.727
Highest quartile (Q4)	1					

OR: odds ratio; CI: confidence interval; BMI: body mass index.